

*J of Immunology 108:253-260, 1972*

AFRRI SR71-7  
MAY 1971

**AFRRI**  
**SCIENTIFIC**  
**REPORT**

**FOR REFERENCE**

Do Not Take From This Room

**ALTERATION IN GRAFT VERSUS HOST  
REACTIVITY AND IN HEMATOPOIETIC  
STEM CELLS OF SPLEEN CELL INOCULA  
FROM DONOR MICE PRETREATED  
WITH PERTUSSIS ANTIGEN**

**ARMED FORCES RADIOBIOLOGY RESEARCH INSTITUTE**  
**Defense Atomic Support Agency**  
**Bethesda, Maryland**

Approved for public release; distribution unlimited


AFRRI SR71-7

All aspects of investigative programs involving the use of laboratory animals sponsored by DoD components are conducted according to the principles enunciated in the "Guide for Laboratory Animal Facilities and Care", prepared by the National Academy of Sciences - National Research Council.


May 1971

ALTERATION IN GRAFT VERSUS HOST REACTIVITY AND IN  
HEMATOPOIETIC STEM CELLS OF SPLEEN CELL INOCULA  
FROM DONOR MICE PRETREATED WITH  
PERTUSSIS ANTIGEN

E. A. EIKMAN  
R. T. BOWSER

  
S. J. BAUM

Chairman  
Experimental Pathology Department

  
HUGH B. MITCHELL  
Colonel, USAF, MC  
Director

ARMED FORCES RADIOBIOLOGY RESEARCH INSTITUTE  
Defense Atomic Support Agency  
Bethesda, Maryland

#### ACKNOWLEDGMENT

The interest and encouragement of S. J. Baum, the technical assistance of R. T. Brandenburg, the statistical analyses performed by S. G. Levin and the editorial assistance of C. H. Poppe are gratefully acknowledged.

# TABLE OF CONTENTS

	Page
Foreword (Nontechnical summary) . . . . .	iii
Abstract . . . . .	v
I. Introduction . . . . .	1
II. Materials and Methods . . . . .	2
Animals . . . . .	2
Radiation . . . . .	2
<u>Bordetella pertussis</u> pretreatment of donor mice . . . . .	2
Cell suspensions . . . . .	3
Assay of GVHR . . . . .	3
Assays of hematopoietic stem cells . . . . .	4
Cytological studies . . . . .	5
Statistical analyses . . . . .	5
III. Results . . . . .	5
IV. Discussion . . . . .	11
Reduction in GVHR . . . . .	11
Increase in indices of hematopoietic stem cells . . . . .	14
Pertussis vaccination modifying hematopoietic transplants . . . . .	15
References . . . . .	17

## LIST OF FIGURES

	Page
Figure 1. Responses of donor mice to pretreatments . . . . .	6
Figure 2. GVHR of C57BL/6J donor spleen cells assayed in B6D2F <sub>1</sub> neonatal recipients . . . . .	7
Figure 3. Indices of hematopoietic stem cells of B6D2F <sub>1</sub> /J donor spleen cells assayed in syngeneic recipients . . . . .	9

## LIST OF TABLES

Table I. GVHR of B6D2F <sub>1</sub> /J Donor Spleen Cells Assayed in Syngeneic Recipients . . . . .	8
Table II. Summary of Effects of Donor Pretreatments on GVHR and Hematopoietic Stem Cells in Spleen Cell Inocula . . . . .	11

## FOREWORD

(Nontechnical summary)

The transplantation of bone marrow from another human is the only specific treatment available to personnel who have received radiation sufficient to cause death from the failure of the blood cell forming (hematopoietic) function of the bone marrow. The transplanted donor cells will generally proliferate in such a host when the host's own marrow is unable to function; a portion of the donor cells, the immunocompetent cells, however, may recognize the host as foreign tissue and mount an immune attack. The outcome is usually fatal. The attack is known as the graft versus host reaction (GVHR); it is analogous to the more familiar host versus graft reaction in which host immunocompetent cells recognize and reject as foreign a transplanted organ. The GVHR may sometimes be reduced if the host and donor tissue types can be matched. Neither this laborious procedure nor other available treatments appear likely to eliminate by themselves the problem of GVHR. We therefore sought a way to provide tissue for transplantation with adequate hematopoietic stem cells but with reduced GVHR. Our studies were designed to test the hypothesis that a dose of pertussis (whooping cough) vaccine administered to donor animals would cause large numbers of immunocompetent lymphocytes to move out of the hematopoietic tissue and into the blood stream. This might reduce the number of GVHR cells in hematopoietic transplants but retain or increase the number of functional hematopoietic stem cells.

Donor mice were treated with either a saline (control) solution or a pertussis vaccine preparation. Donor spleen cells, which in the mouse resemble human bone

marrow in their functions, were then transplanted by injection into recipient mice in experiments designed to test GVHR and hematopoietic stem cells quantitatively and specifically.

We observed that pertussis vaccination reduced the proportion of GVHR cells and increased the proportion of hematopoietic stem cells in donor mouse spleens. These two effects improved the treatment value of this hematopoietic transplant thirteenfold compared with control transplants. The findings suggest a general approach to GVHR which may have implications for the use of hematopoietic transplantation in cancer therapy, some anemias, and other conditions in addition to radiation injury. More animal experiments will be necessary to clarify the mechanism of the effects observed and to evaluate the clinical implications these results may have.



## ABSTRACT

The graft versus host reactivity (GVHR) and hematopoietic stem cells of spleen cell inocula were assayed 3 days after pretreatment of donor mice with heated or unheated pertussis vaccine. Unheated vaccine reduced GVHR to 17 percent of control reactivity, measured by the Simonsen spleen weight assay, and increased 2.2-fold the proportion of hematopoietic stem cells, assayed as colony forming units (CFU). Heated vaccine, which did not produce the lymphocytosis induced in donors by unheated vaccine, reduced GVHR to 38 percent of control reactivity, and increased the proportion of CFU 3.8-fold. Alternative mechanisms for these effects are discussed.

## I. INTRODUCTION

Allogeneic bone marrow transplants prolong survival in rodents with radiation-induced hematopoietic aplasia<sup>20,26</sup> through establishment of hematopoietic chimerism by the transplant.<sup>2,28</sup> However, in humans with hematopoietic aplasia, clinical trials of bone marrow transplantation demonstrated that the establishment of a histoincompatible marrow allograft leads to an acute fatal secondary syndrome.<sup>30,31,51</sup> The syndrome is attributed to an immunologic graft versus host reaction (GVHR).<sup>42,45,52</sup> Recently introduced histocompatibility matching methods offer hope of alleviating the syndrome when suitable donors are available.<sup>1,8,15,18</sup> Additional methods of reducing the GVHR are required especially for the majority of recipients for whom matched donors may not be available. Several methods of reducing the GVHR are under investigation in other laboratories and have been recently summarized.<sup>10</sup>

Pertussis vaccine mobilizes mouse lymphocytes<sup>38</sup> of a thymus-dependent population<sup>21</sup> from the hematopoietic organs into the blood stream. Thymus-dependent lymphocytes may initiate cellular immune reactions<sup>14,17,37</sup> of which the GVHR is an example.<sup>50</sup> We therefore hypothesized that pertussis vaccine pretreatment of donor animals would deplete donor organs of GVHR cells while retaining significant hematopoietic stem cells. Transplantation of mouse spleen cells from pretreated donors was chosen as an experimental model because mouse spleen resembles human bone marrow in containing both hematopoietic stem cells and cells causing acute GVHR.<sup>40</sup> The studies reported in this communication showed that pertussis vaccination of donor mice leads to a reduction of GVHR and to an increase in hematopoietic stem cells of donor mouse spleen. It was also shown that lymphocyte mobilization could only partially account for these effects.

## II. MATERIALS AND METHODS

Animals. C57BL/6J ♀ (H-2<sup>b</sup>), DBA/2J ♂ (H-2<sup>d</sup>), and the F<sub>1</sub> ♀ hybrid offspring of these strains of mice (designated B6D2F<sub>1</sub>/J) were purchased from the Jackson Laboratory, Bar Harbor, Maine. The animals were allowed food,\* and water adjusted to pH 3 with hydrochloric acid, ad libitum. Donors and irradiated recipients were used at 9-12 weeks of age, weighing 17-24 g. Neonatal litters of B6D2F<sub>1</sub> mice were bred in our laboratory from Jackson stock; they were used in assays for GVHR at 6-10 days of age and were nursed by the mother until completion of the assays.

Radiation. Irradiated animals received 850 rads (LD<sub>95-100/30</sub>) produced by a General Electric Maxitron x-ray machine with the following characteristics: 250 kVp, 30 mA, 1.2 mm Be + 0.95 mm Cu filtration (HVL = 1.91 mm Cu), target to subject midline distance 67.5 cm. The midline absorbed dose rate was 56.2 rads/minute.

Bordetella pertussis pretreatment of donor mice. B. pertussis vaccine† containing  $200 \times 10^9$  killed organisms/ml (6 mg protein/ml)‡ in 0.01 percent thimerosal was diluted to 5 volumes with 0.9 percent sodium chloride just prior to use. Heated pertussis vaccine was prepared by heating diluted vaccine at 100°C for 30 minutes. Heating in this manner removes the capacity of the vaccine to mobilize lymphocytes;<sup>38</sup> it also removes immunogenicity for specific protection from pertussis infection, and a number of other activities of the vaccine.<sup>23</sup> Ten billion organisms contained in 0.25 ml of the unheated or heated diluted vaccine were injected into a tail vein of

---

\* D & G Rat and Mouse Diet, specific pathogen free. G. L. Baking Company, Frederick, Maryland

† Lot No. 985739, kindly supplied by Dr. H. B. Devlin, Parke, Davis and Company, Detroit, Michigan

‡ Kindly determined by Dr. W. D. Skidmore by the method of Lowry et al.<sup>27</sup>

experimental donors; control donors received 0.25 ml saline. Three days after injection, the animals were sacrificed by cervical dislocation and the spleens were removed and weighed. The number of tail vein blood leukocytes of about one-half the donor animals were counted as an index of stability of the vaccine used. The potency of the vaccine for producing leukocytosis or increased spleen weights was stable during the course of the studies.

Cell suspensions. The spleens removed from donor animals were weighed and maintained subsequently in Hanks' balanced salt solution (HBSS) in an ice bath. The spleens were minced with scissors and expressed through 400-mesh nylon gauze. The resulting suspension of individual cells was washed by dilution in 30 volumes of HBSS followed by centrifugation at 330 g for 8 minutes at 4°C. The supernatant was then decanted; this procedure was repeated for a total of three washes. The cells were then resuspended with HBSS to about 0.3 ml/spleen. An aliquot was diluted in a red cell pipet with 0.4 percent trypan blue in saline. Nucleated cells excluding trypan blue were counted in a hemacytometer 1-3 minutes after dilution. The suspensions were then diluted with HBSS to contain the desired number of nucleated cells either in 0.4 ml for tail vein injection into adult irradiated recipients or in 0.1 ml for intra-peritoneal injection into neonates.

Assay of GVHR. A modification of the spleen weight assay of Simonsen<sup>34, 46</sup> was performed. The assay is based on the early finding of splenomegaly in neonatal mice following injection of histoincompatible adult lymphoid cells. Doses of 2, 5, 10 or 20 million spleen cells from experimental or control donors were injected intra-peritoneally into 6- to 10-day-old B6D2F<sub>1</sub> mice. Three or four littermates were left

uninjected. Litters of fewer than six animals were not used and animals over 10 in an individual litter were discarded. After the injection, the litter was returned to the mother for 9 days. The litter was then sacrificed and individual spleen and body weights were determined. The spleen index for each of the injected animals was calculated by dividing its spleen to body weight ratio by the average of the spleen to body weight ratios of the uninjected littermates.

Assays of hematopoietic stem cells. The hematopoietic stem cells of experimental or control donor spleen cell suspensions were determined by intravenous injections of  $2.0 \times 10^5$  or  $1.0 \times 10^6$  cells into lethally irradiated (850 rads) syngeneic recipients. Irradiated control recipients received 0.4 ml HBSS. Untreated unirradiated control mice were also maintained and subjected to the hematopoietic assays. On the 10th day, 1  $\mu$ Ci of  $^{59}\text{Fe}^*$  was injected subcutaneously.<sup>29</sup> Six hours later the mice were sacrificed and spleen and body weights determined. The radioactivity of a humerus and the spleen was determined in a scintillation counter. The spleens were fixed in Bouin's solution and the number of macroscopic surface nodules on the spleen, each representing a hematopoietic colony forming unit (CFU), were counted.<sup>49</sup>

In preliminary studies, the hematopoietic assays of C57BL/6J mice used as donors in the GVHR assay were compared with those of B6D2F<sub>1</sub>/J mice. Since spleen cells of pertussis vaccine pretreated or control donors from both strains produced similar hematopoietic responses in syngeneic recipients, all subsequent studies reported here were performed with B6D2F<sub>1</sub>/J donors in order to avoid the high early postirradiation mortality of C57BL/6 mice.<sup>32</sup>

---

\* As ferrous citrate at pH 6.8. New England Nuclear Corporation, Boston, Massachusetts

Cytological studies. In separate experiments, animals of each of the donor strains were injected with either unheated pertussis vaccine, heated pertussis vaccine or saline as described above. At 3 days, total and differential leukocyte counts were determined, body and spleen weights were measured, and a touch preparation of the spleen was stained with Wright-Leishman's stain. Total nucleated cells per spleen were calculated from counts of aliquots of pooled spleen suspensions diluted in 3 per-cent acetic acid.

Statistical analyses. The t-test was used to compare averages of values obtained. Differences are reported as significant when  $p < 0.05$ .

### III. RESULTS

Effects of the vaccine preparations used on donor mice are shown in Figure 1. Both the heated and unheated pertussis vaccine caused a significant increase over values from control animals in average spleen weight and in the number of nucleated cells per spleen. Body weights were not significantly affected. Examination of the tail vein blood showed approximately a ninefold increase in peripheral leukocytes, with an eightfold increase in the sum of lymphocytes and monocytes in the animals treated with unheated vaccine. The heated vaccine did not significantly change the blood counts.

Cytological examination of touch preparations of the spleen showed increases over controls in immature cells of the myelocytic and lymphocytic series and in the numbers of mature neutrophils in the spleens of the animals treated with heated or unheated vaccine.




Figure 1. Responses of donor mice to pretreatments. Average values for four or five recipients are shown 3 days after injection of heated or unheated pertussis vaccine or saline solution. The vertical bar ( $\top$ ) denotes 1 S. E. except in the case of spleen cells where 1 S. D. for four replicate measurements from pooled spleens is shown.

Results of the assay of the GVHR are shown in Figure 2 for spleen cells of C57BL/6J donors from each of the pretreatment groups injected into B6D2F<sub>1</sub> recipients. Compared with the control curve, both of the experimental dose response curves indicate that more experimental cells are required to produce a given spleen



index. Statistical comparison of the slopes of the three regression lines<sup>16</sup> shows that they are not significantly different. The relative reactivities of the various cell populations may therefore be expressed as the ratio of the numbers of control cells to experimental cells required to produce any spleen index within the range of indices associated with parallel dose response curves. The ratio may be obtained through an equation derived for calculation of the distance between regression lines.<sup>16</sup> Substituting the GVHR data shown in Figure 2 into the equation, one obtains the ratios of 0.38 for the numbers of control cells to cells from donors pretreated with heated vaccine, and 0.17 for the numbers of control cells to cells from donors pretreated with unheated vaccine.




Figure 2. GVHR of C57BL/6J donor spleen cells assayed in B6D2F<sub>1</sub> neonatal recipients. Cells from control donors pretreated with saline or from experimental donors pretreated with either heated or unheated pertussis vaccine were injected into B6D2F<sub>1</sub> neonates in graduated doses in six experiments. Each point represents the average spleen index  $\pm$  1 S. E. of a total of 30-40 recipients. Regression lines were fitted by the method of least squares.

As a control experiment for the assay of the GVHR, separate assays reported in Table I were performed in B6D2F<sub>1</sub> neonates but, instead of the parental strain, donors were syngeneic B6D2F<sub>1</sub>/J adults receiving pretreatment identical to that of the



C57BL/6J parental strain donors. The observed absence of significant changes in the average spleen indices of neonatal B6D2F<sub>1</sub> recipients after injection with syngeneic cells from control or experimental donors is consistent with the interpretation that the splenomegaly induced in the spleen assay of the GVHR required a histocompatibility difference between donor and recipient, and excludes the possibility that the GVHR assay shown in Figure 2 was influenced by factitious splenomegaly such as might be caused by vaccine carried with the donor cells.

Table I. GVHR of B6D2F<sub>1</sub>/J Donor Spleen Cells Assayed in Syngeneic Recipients

Donor pretreatment (I. V.)	Recipient cell dose (millions)	Number of recipients	Average spleen index $\pm$ 1 S. E.
Saline	2	6	$1.00 \pm 0.07^*$
	5	5	$1.14 \pm 0.08$
	10	5	$1.00 \pm 0.09$
Heated pertussis vaccine	5	5	$0.99 \pm 0.10$
	10	6	$0.99 \pm 0.02$
	20	6	$1.08 \pm 0.05$
Unheated pertussis vaccine	5	5	$0.97 \pm 0.07$
	10	6	$0.97 \pm 0.08$
	20	6	$1.12 \pm 0.08$

\* None of the averages differ significantly from 1.0

Hematopoietic stem cells of B6D2F<sub>1</sub>/J spleen cells from control and experimental donors are shown in Figure 3. Although the standard errors of the average indices in these experiments were relatively large, the spleen colony counts, splenic iron uptake and spleen weight assays (Figure 3A-C) were in agreement when experimental inocula were compared with control cell inocula. Inocula from donors pretreated with heated




Figure 3. Indices of hematopoietic stem cells of B6D2F<sub>1</sub>/J donor spleen cells assayed in syngeneic recipients. Recipients were lethally irradiated, then injected with HBSS or with spleen cell suspensions from control donors pretreated with saline or from experimental donors pretreated with either heated or unheated pertussis vaccine. From 19-23 survivors in each group were assayed on the 10th day for (A) spleen colonies, (B) spleen  $^{59}\text{Fe}$  uptake, (C) spleen weight, (D) humerus  $^{59}\text{Fe}$  uptake, and (E) body weight.

vaccine produced the largest average indices, followed by inocula from donors pretreated with unheated vaccine. Control cell inocula produced the smallest indices. Inocula from donors pretreated with heated vaccine resulted in increased average body weights in the recipients but inocula from donors pretreated with unheated vaccine resulted in average body weights about the same as those resulting from control cell inocula.

Since the number of CFU counted is linearly related to the number of hematopoietic stem cells injected<sup>49</sup> up to a saturation point of about 25-30 CFU per spleen, quantitative comparisons of the stem cell content of the various inocula may be drawn. Thus, when effects at a dose of 0.2 million cells are compared, spleen cells from donors pretreated with unheated or heated pertussis vaccine produce respectively 2.2 or 3.8 times as many CFU as the cells from control donors (Figure 3A).

The determinations of splenic iron uptake were controlled by assessment of marrow iron uptake (Figure 3D). Marrow erythropoiesis, as reflected in the uptake by the humerus, remained at the reduced level of the irradiated mice which received HBSS without cells. The lack of evidence of change in marrow iron uptake under any of the experimental conditions employed excludes the possibility that differences in recipient spleen erythropoiesis observed were merely reflecting proportionate changes in marrow erythropoiesis without actual changes in the total effective erythropoietic stem cells transplanted.

The combined effects of the vaccine pretreatments on donor GVHR and hematopoietic stem cells may be summarized in the manner of Dicke et al.<sup>9</sup> as the CFU to GVHR ratio. The ratios shown in Table II represent an approximate index of the

therapeutic efficacy of the donor spleen cells in hematopoietic transplantation. A ratio of 13 was found for unheated pertussis vaccine pretreatment and 10 for heated pertussis vaccine pretreatment, compared with the control index of 1.

Donor pretreatment (I. V.)	Fraction of control index		Ratio* CFU to GVHR
	CFU	GVHR	
Heated pertussis vaccine	3.8	0.38	10
Unheated pertussis vaccine	2.2	0.17	13

Table II. Summary of Effects of Donor Pretreatments on GVHR and Hematopoietic Stem Cells in Spleen Cell Inocula

\* Ratio is 1 for control inoculum

#### IV. DISCUSSION

The present studies showed that following pertussis vaccination, the GVHR of donor mouse spleen cells was reduced to 17 percent of control reactivity when the cells were transplanted across an H-2 histocompatibility difference. Pertussis vaccination was also shown to lead to more than twice the number of hematopoietic stem cells in donor spleens when assayed as splenic CFU in irradiated syngeneic recipients. Each of these findings will be discussed separately, followed by a discussion of their implications when considered together.

Reduction in GVHR. Other evidence had suggested that the increase in blood lymphocyte counts following pertussis vaccination results through mobilization of thymus-dependent lymphocytes,<sup>21, 38</sup> the cell population thought to cause the GVHR.<sup>14, 17, 37</sup> The vaccine used in the present studies increased donor blood lymphocyte counts, and also reduced the proportion of GVHR cells in donor spleens. When the vaccine was heated to remove its capacity to mobilize lymphocytes, however, it was still effective in reducing the GVHR to 38 percent of control reactivity.

Therefore, not all of the effect of the pertussis vaccine on the GVHR is likely to be due to the mobilization of lymphocytes.

If mobilization of the GVHR cells into the peripheral blood is important in the effect of the unheated vaccine in reducing the GVHR of the donor spleen, one would predict that the net GVHR of the peripheral blood would be increased. Prior comparisons of the proportion of GVHR cells of peripheral blood from pertussis vaccine pretreated and control donors showed decreased reactivity in the mouse<sup>6</sup> but increased reactivity in the rat.<sup>24</sup> To quantitate the total blood GVHR relative to control or other organ reactivity, however, it would be necessary to know the volume of distribution of the reactive cells in the donor and to use a GVHR assay demonstrated to produce effects dependent upon the number of reactive cells injected. In neither of the studies of peripheral blood were these requirements met. Thus, it remains to be answered whether there is an increase in the total number of GVHR cells in the peripheral blood after pertussis vaccination.

Since we have no direct proof that unheated vaccine shifts GVHR cells into the blood, and since the heated vaccine reduced spleen GVHR without apparently mobilizing cells, mechanisms other than lymphocyte mobilization should be considered to account at least partially for the ability of the vaccine to reduce the GVHR. A simple alternative mechanism would be the dilution in the donor spleen of GVHR cells by unreactive cells. This would lead to a reduction in the proportion of GVHR cells.

Unheated vaccine led to a 2.1-fold increase in the number of spleen cells, which could account for a reduction to as little as 48 percent of control reactivity; heated vaccine led to a 1.6-fold increase in the number of spleen cells, which could account

for a reduction to about 63 percent of control reactivity. Thus, dilution alone could account for a major portion of the observed reduction in GVHR if a large portion of the additional cells in the spleens were inactive in the assay.

A third possible mechanism for the observed reduction in the GVHR is reduced entry of the parental spleen GVHR cells into the spleens of the  $F_1$  neonatal recipients. Recipient splenomegaly and thus apparent GVHR would thereby be diminished. This may be unlikely if the behavior of our spleen cells is comparable to that reported by Morse and Barron for labeled lymphocytes from the peripheral blood of animals undergoing pertussis vaccine-induced lymphocytosis. Compared with controls there was no reported difference in the entry of these peripheral blood cells into the spleens of adult syngeneic recipients.<sup>39</sup>

A fourth possibility is that pertussis vaccine reduces the GVHR by inactivating GVHR cells through antigenic competition with recipient histocompatibility antigens. The reduction of GVHR by a variety of other antigens under a variety of conditions has been attributed to antigenic competition.<sup>19,25,36</sup> Antigenic competition might also explain the finding that under some conditions guinea pigs receiving pertussis vaccine have shown diminished tuberculin reactivity<sup>12</sup> and the finding that fowl receiving pertussis vaccine showed diminished cutaneous lymphocyte transfer reactions.<sup>43</sup> The precise mechanism of antigenic competition in cellular immune responses is, however, not clear.

In contrast to the immunosuppressive effects of pertussis vaccine in reducing the GVHR, the vaccine is active as an adjuvant in humoral immune responses.<sup>11,41</sup> Under conditions comparable to ours, Cantor et al.<sup>5</sup> showed pretreatment by the



adjuvant polyinosinic-polycytidylic acid given to donor BALB/c mice increased the GVHR of donor spleen cells. Similarly, an adjuvant effect of the pertussis vaccine in cellular immunity would have been expected to lead to an increase rather than a decrease in the GVHR under the conditions of the present experiments. Either our vaccine was not effective as an adjuvant under the present conditions or a quantitatively greater immunosuppressive effect may have obscured its adjuvanticity.

Increase in indices of hematopoietic stem cells. Pretreatment of donor mice with unheated pertussis vaccine more than doubled the proportion of hematopoietic stem cells in the spleen cell inoculum, but pretreatment with heated vaccine more than tripled this proportion. The difference in the activities of the two preparations could be accounted for if, as in the case of lymphocyte mobilization, the unheated vaccine mobilizes hematopoietic stem cells and the heated vaccine does not. Given equal stimulation of hematopoietic stem cells, the spleens from the animals in which the cells were not mobilized would contain more stem cells. This hypothesis is in accord with prior findings that unheated vaccine leads to an increase in the circulating hematopoietic stem cells of mice.<sup>3,6</sup> The effect of heated vaccine on circulating stem cells was not reported.

The finding that donor spleen hematopoietic stem cells are increased after pretreatment with either heated or unheated vaccine adds support to the interpretation that endotoxin, which is heat stable,<sup>22</sup> may be responsible for the increase in hematopoiesis which pertussis vaccine may produce.<sup>13</sup> Other bacterial endotoxin preparations<sup>47,48</sup> as well as a variety of other antigens,<sup>4,33</sup> stimulate erythropoiesis. Studies using purified B. pertussis endotoxin might clarify this possible relationship.

The observation that recipient bone marrow iron uptake remained depressed to the level of the irradiated controls regardless of the treatment or dose employed is in accord with findings by others that marrow erythropoiesis is not an important component of the erythropoietic recovery of these irradiated mice 10 days after radiation and transplantation of hematopoietic tissue.<sup>35</sup>

Pertussis vaccination modifying hematopoietic transplants. A theoretical increase in the efficacy of the transplantation of donor spleen cells is suggested by the CFU to GVHR ratio of 13 times that of the control cells. Since reduction in the GVHR may synergistically increase hematopoiesis,<sup>7, 44</sup> the actual improvement in therapeutic efficacy in allogeneic transplantation may be even greater.

Parallel quantitative measurements of the GVHR and hematopoietic stem cells allow at least approximate comparison of alternative methods of preparing hematopoietic tissue for transplantation. For example, in the studies of Dicke et al.,<sup>9</sup> mouse spleen cell fractionation on a discontinuous albumin gradient produced a fraction with more than a tenfold increase in hematopoietic cells and less than 10 percent of the GVHR compared to the control suspension. The resulting CFU to GVHR ratio of greater than 100 for the 3-5 percent of cells recovered may be compared with the CFU to GVHR ratio of 13 found in the present study.

The more general use by other investigators of parallel quantitative assays of the GVHR and hematopoiesis would aid in further comparisons of treatments and would facilitate the evaluation of two or more treatment methods applied to the same hematopoietic transplant.



Further studies seem warranted to determine whether pertussis vaccine, its modified fractions, or other agents which duplicate useful properties of the vaccine, may have eventual therapeutic implications in human marrow transplantation.

## REFERENCES

1. Bach, F. H., Albertini, R. J., Joo, P., Anderson, J. L. and Bortin, M. M. Bone-marrow transplantation in a patient with the Wiskott-Aldrich syndrome. *Lancet* 2:1364-1366, 1968.
2. Barnes, D. W. H. and Loutit, J. F. Spleen protection: the cellular hypothesis. In: Radiobiology Symposium, Proceedings of the Symposium held at Liège, August - September 1954, Bacq, Z. M. and Alexander, P., editors, pp. 134-135. New York, Academic Press; London, Butterworths Scientific Publications, 1955.
3. Barnes, D. W. H. and Loutit, J. F. Effects of irradiation and antigenic stimulation on circulating haemopoietic stem cells of the mouse. *Nature* 213:1142-1143, 1967.
4. Boggs, D. R., Marsh, J. C., Chervenick, P. A., Bishop, C. R., Cartwright, G. E. and Wintrobe, M. M. Factors influencing hematopoietic spleen colony formation in irradiated mice. II. The effect of foreign materials. *J. Exptl. Med.* 126:851-870, 1967.
5. Cantor, H., Asofsky, R. and Levy, H. B. The effect of polyinosinic-polycytidylic acid upon graft-vs-host activity in BALB/c mice. *J. Immunology* 104:1035-1038, 1970.
6. Cole, L. J. and Maki, S. E. Hemopoietic colony-forming cells in mouse blood: Effect of pertussis vaccine injection and of thymectomy-irradiation. *Exptl. Hematology* 15:82-85, 1968.
7. Davis, W. E. and Cole, L. J. Suppression of hemopoietic colony-forming units in mice by graft-versus-host reactions. *Transplantation* 5:60-67, 1967.
8. De Koning, J., van Bekkum, D. W., Dicke, K. A., Dooren, L. J., van Rood, J. J. and Rádl, J. Transplantation of bone-marrow cells and fetal thymus in an infant with lymphopenic immunological deficiency. *Lancet* 1:1223-1227, 1969.
9. Dicke, K. A., van Hooft, J. I. M. and van Bekkum, D. W. The selective elimination of immunologically competent cells from bone marrow and lymphatic cell mixtures. *Transplantation* 6:562-570, 1968.
10. Fahey, J. L., Mann, D. L., Asofsky, R. and Rogentine, G. N. Recent progress in human transplantation immunology. *Ann. Intern. Med.* 71:1177-1196, 1969.
11. Farthing, J. R. The role of Bordetella pertussis as an adjuvant to antibody production. *Brit. J. Exptl. Pathol.* 42:614-622, 1961.

12. Floersheim, G. L. Effect of pertussis vaccine on the tuberculin reaction. *Int. Arch. Allergy* 26:340-344, 1965.
13. Fruhman, G. J. Hematopoietic changes in mice following Bordetella pertussis vaccine. *Proc. Soc. Exptl. Biol. Med.* 124:728-730, 1967.
14. Gabrielsen, A. E., Cooper, M. D., Peterson, R. D. A. and Good, R. A. The primary immunologic deficiency diseases. In: *Textbook of Immunopathology*, Vol. II, p. 385, Miescher, P. A. and Müller-Eberhard, H. J., editors. New York and London, Grune and Stratton, 1969.
15. Gatti, R. A., Meuwissen, H. J., Allen, H. D., Hong, R. and Good, R. A. Immunological reconstitution of sex-linked lymphopenic immunological deficiency. *Lancet* 2:1366-1369, 1968.
16. Geigy Chemical Corporation. Statistical methods, p. 177. In: *Documenta Geigy Scientific Tables*, 6th ed., Diem, K., editor. Ardsley, New York, Geigy Pharmaceuticals, 1962.
17. Gowans, J. L. and McGregor, D. D. The immunological activities of lymphocytes. *Progr. Allergy* 9:1-78, 1965.
18. Hitzig, W. H., Kay, H. E. M. and Cottier, H. Familial lymphopenia with agammaglobulinaemia: An attempt at treatment by implantation of foetal thymus. *Lancet* 2:151-154, 1965.
19. Hunter, R. L., Millman, R. B. and Lerner, E. M., II. Suppression of graft-versus-host disease and antibody formation by phytohemagglutinin. *Transplantation* 8:413-421, 1969.
20. Jacobson, L. O., Marks, E. K., Robson, M. J., Gaston, E. and Zirkle, R. E. The effect of spleen protection on mortality following x-irradiation. *J. Lab. Clin. Med.* 34:1538-1543, 1949.
21. Kalpaktsoglou, P. K., Yunis, E. J. and Good, R. A. Changes produced by pertussis antigen on the blood cells and lympho-haemopoietic tissues after early and late thymectomy or splenectomy. *Clin. Exptl. Immunol.* 5:91-103, 1969.
22. Kind, L. S. The altered reactivity of mice after inoculation with Bordetella pertussis vaccine. *Bacteriol. Rev.* 22:173-182, 1958.
23. Levine, L. and Pieroni, R. E. A unitarian hypothesis of altered reactivity to stress mediated by Bordetella pertussis. *Experientia* 22:797-798, 1966.

24. Levine, S. and Iwahara, M. Graft-versus-host disease produced with whole blood. Enhancement by pertussis vaccination of donors. *Transplantation* 8: 462-465, 1969.
25. Liacopoulos, P., Merchant, B. and Harrell, B. E. Inhibition of the graft-versus-host reaction by pretreatment of donors with various antigens. *Transplantation* 5:1423-1435, 1967.
26. Lorenz, E., Uphoff, D., Reid, T. R. and Shelton, E. Modification of irradiation injury in mice and guinea pigs by bone marrow injections. *J. Nat. Cancer Inst.* 12:197-201, 1951.
27. Lowry, O. H., Rosebrough, N. J., Farr, A. L. and Randall, R. J. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* 193:265-275, 1951.
28. Main, J. M. and Prehn, R. T. Successful skin homografts after administration of high dosage X radiation and homologous bone marrow. *J. Nat. Cancer Inst.* 15:1023-1029, 1955.
29. Marsh, J. C., Boggs, D. R., Bishop, C. R., Chervenick, P. A., Cartwright, G. E. and Wintrobe, M. M. Factors influencing hematopoietic spleen colony formation in irradiated mice. I. The normal pattern of endogenous colony formation. *J. Exptl. Med.* 126:833-849, 1967.
30. Mathé, G., Amiel, J. L., Schwarzenberg, L., Schneider, M., Cotton, A., Schlumberger, J. R., Nouza, K. and Hrask, Y. Bone marrow transplantation in man. *Transplantation Proc.* 1:16-24, 1969.
31. Mathé, G., Bernard, J., de Vries, M. J., Schwarzenberg, L., Larrieu, M. J., Lallanne, C. M., Dutreix, A., Amiel, J. L. and Surmont, J. Nouveaux essais de greffe de moelle osseuse homologue après irradiation totale chez des enfants atteints de leucémie aiguë en rémission. Le problème du syndrome secondaire chez l'homme. *Rev. Hématol. (Paris)* 15:115-161, 1960.
32. McCulloch, E. A. and Till, J. E. Repression of colony-forming ability of C57BL hematopoietic cells transplanted into non-isologous hosts. *J. Cellular Comp. Physiol.* 61:301-308, 1963.
33. Metcalf, D. Studies on colony formation in vitro by mouse bone marrow cells. I. Continuous cluster formation and relation of clusters to colonies. *J. Cellular Physiol.* 74:323-332, 1969.
34. Michie, D. The biometrics of the spleen weight assay. In: *Handbook of Experimental Immunology*, Weir, D. M., editor, pp. 969-987. Philadelphia, Pennsylvania, F. A. Davis Company, 1967.

35. Micklem, H. S. and Loutit, J. F. Tissue Grafting and Radiation, p. 78. New York and London, Academic Press, 1966.
36. Miller, J., Martinez, C. and Good, R. A. Reciprocal competition of a variety of antigens in the suppression of immunologic reactivity. J. Immunology 93: 342-351, 1964.
37. Miller, J. F. A. P. and Osoba, D. Current concepts of the immunological function of the thymus. Physiol. Rev. 47:437-520, 1967.
38. Morse, S. I. Studies on the lymphocytosis induced in mice by Bordetella pertussis. J. Exptl. Med. 121:49-68, 1965.
39. Morse, S. I. and Barron, B. A. Studies on the leukocytosis and lymphocytosis induced by Bordetella pertussis. III. The distribution of transfused lymphocytes in pertussis-treated and normal mice. J. Exptl. Med. 132:663-672, 1970.
40. Muller-Bérat, C. N., van Putten, L. M. and van Bekkum, D. W. Cytostatic drugs in the treatment of secondary disease following homologous bone marrow transplantation: Extrapolation from the mouse to the primate. Ann. N. Y. Acad. Sci. 129:340-354, 1966.
41. Pieroni, R. E. and Levine, L. Adjuvant principle of pertussis vaccine in the mouse. Nature 211:1419-1420, 1966.
42. Schwartz, E. E., Upton, A. C. and Congdon, C. C. A fatal reaction caused by implantation of adult parental spleen tissue in irradiated F<sub>1</sub> mice. Proc. Soc. Exptl. Biol. Med. 96:797-800, 1957.
43. Seiler, K. Prüfung von Antiphlogistica an einer immunologischen Entzündung des Hühnchens. Arch. Int. Pharmacodyn. 169:452-468, 1967.
44. Sensenbrenner, L. L. Simultaneous assay of the effects of agents on GvH disease and hematopoietic spleen colony forming units. Exptl. Hematology (in press).
45. Simonsen, M. The impact on the developing embryo and newborn animal of adult homologous cells. Acta Pathol. Microbiol. Scand. 40:480-500, 1957.
46. Simonsen, M. Graft versus host reactions. Their natural history and applicability as tools of research. Progr. Allergy 6:349-467, 1962.
47. Smith, W. W., Alderman, I. M. and Gillespie, R. E. Increased survival in irradiated animals treated with bacterial endotoxins. Am. J. Physiol. 191: 124-130, 1957.

48. Smith, W. W., Marston, R. Q. and Cornfield, J. Patterns of hemopoietic recovery in irradiated mice. *Blood* 14:737-747, 1959.
49. Till, J. E. and McCulloch, E. A. A direct measurement of the radiation sensitivity of normal mouse bone marrow cells. *Radiation Res.* 14:213-222, 1961.
50. Trentin, J. J. and Session, J. Immunological tolerance and lymphoid chimerism following injection of homologous spleen cells into newborn mice. In: *Mechanisms of Immunological Tolerance, Proceedings of Symposium held at Liblice near Prague, November 8-10, 1961*, Hasěk, M., Lengerová, A. and Vojtišková, M., editors, pp. 31-34. Prague, Publishing House of the Czechoslovak Academy of Sciences, 1962.
51. van Bekkum, D. W. Hostile grafts. In: *Advance in Transplantation*, Dausset, J., Hamburger, J. and Mathé, G., editors, p. 565. Baltimore, Maryland. The Williams and Wilkins Company, 1968.
52. van Bekkum, D. W. and de Vries, M. J. *Radiation Chimaeras*, Chapter III. London, Logos Press Limited, 1967.



## DISTRIBUTION LIST

### AIR FORCE

The Surgeon General, U. S. Department of the Air Force, Washington, D. C. 20333 (1)  
Executive Officer, Director of Professional Services, Office of the Surgeon General, Hq. USAF (AFMSPA) T-8,  
Washington, D. C. 20333 (1)  
Headquarters, U. S. Air Force (AFMSPAB), Washington, D. C. 20333 (1)  
USAFSAM (SMBR), ATTN: Chief, Radiobiology Branch, Brooks AFB, Texas 78235 (1)  
Air Force Weapons Laboratory, ATTN: WLIL (1), ATTN: WLRB-2 (1), Kirtland AFB, New Mexico 87117 (2)  
Chief, Nuclear Medicine Department, P. O. Box 5088, USAF Hospital, Wright-Patterson AFB, Ohio 45433 (1)  
Office of the Command Surgeon (ADCSG), Hq. ADC, USAF, Ent AFB, Colorado 80912 (1)  
Commander, 6571st Aeromedical Research Laboratory, Holloman AFB, New Mexico 88330 (2)

### ARMY

The Surgeon General, U. S. Department of the Army, Washington, D. C. 20315 (1)  
Surgeon General, ATTN: MEDDH-N, U. S. Department of the Army, Washington, D. C. 20315 (1)  
USACDC CSSG, Doctrine Division, Fort Lee, Virginia 23801 (1)  
CG, USCONARC, ATTN: ATUTR-TNG (NBC), Fort Monroe, Virginia 23351 (1)  
Commanding Officer, U. S. Army Medical Research Laboratory, Fort Knox, Kentucky 40121 (1)  
Commanding Officer, USA Nuclear Medical Research Detachment, Europe, APO New York, N. Y. 09180 (2)  
Army Research Office, ATTN: Chief, Scientific Analysis Branch, Life Sciences Division, 3045 Columbia Pike,  
Arlington, Virginia 22204 (1)  
Division of Nuclear Medicine, Walter Reed Army Institute of Research, Walter Reed Army Medical Center  
Washington, D. C. 20012 (5)  
Commanding Officer, U. S. Army Environmental Hygiene Agency, ATTN: USAEHA-RP, Edgewood Arsenal, Mary-  
land 21010 (1)  
Commandant, U. S. Army Medical Field Service School, ATTN: MEDEW-ZNW, Fort Sam Houston, Texas 78234 (1)

### NAVY

Chief, Bureau of Medicine and Surgery, U. S. Navy Department, Washington, D. C. 20390 (1)  
Chief, Bureau of Medicine and Surgery, ATTN: Code 71, U. S. Navy Department, Washington, D. C. 20390 (1)  
Director, Biological Sciences Division, Office of Naval Research, Washington, D. C. 20360 (1)  
Commanding Officer, Naval Aerospace Medical Institute, NAMC, ATTN: Research Director, Pensacola, Florida  
32512 (3)  
Head, Animal Behavioral Sciences Branch, Naval Aerospace Medical Institute, Naval Aerospace Medical Center,  
ATTN: Dr. John S. Thach, Jr., Pensacola, Florida 32512 (1)  
Commanding Officer, U. S. Naval Hospital, ATTN: Director, REEL, NNMC, Bethesda, Maryland 20014 (1)  
Commanding Officer, Nuclear Weapons Training Center, Atlantic, Nuclear Warfare Department, Norfolk, Virginia  
23511 (1)  
Commanding Officer, Naval Submarine Medical Center, Naval Submarine Base, NL, ATTN: Medical Library,  
Groton, Connecticut 06340 (1)  
Commanding Officer, Naval Submarine Medical Center, Naval Submarine Base, NL, ATTN: Code 53, Nuclear Medi-  
cine Training Division, Groton, Connecticut 06340 (1)

### D. O. D.

Director, Defense Atomic Support Agency, Washington, D. C. 20305 (1)  
Director, Defense Atomic Support Agency, ATTN: DDST, Washington, D. C. 20305 (1)  
Director, Defense Atomic Support Agency, ATTN: Chief, Medical Directorate, Washington, D. C. 20305 (4)  
Director, Defense Atomic Support Agency, ATTN: Technical Library (APTL), Washington, D. C. 20305 (2)  
Commander, Field Command, Defense Atomic Support Agency, ATTN: FC Technical Library, Sandia Base,  
Albuquerque, New Mexico 87115 (1)  
Director, Armed Forces Institute of Pathology, Washington, D. C. 20305 (1)  
Administrator, Defense Documentation Center, Cameron Station, Bldg. 5, Alexandria, Virginia 22314 (20)

### OTHER GOVERNMENT

U. S. Atomic Energy Commission, Headquarters Library, Reports Section, Mail Station G-17, Washington, D. C.  
20545 (1)  
U. S. Atomic Energy Commission, Division of Biology and Medicine, Washington, D. C. 20545 (1)  
U. S. Atomic Energy Commission, Bethesda Technical Library, 7920 Norfolk Avenue, Bethesda, Maryland 20014 (1)

## OTHER GOVERNMENT (continued)

- National Aeronautics and Space Administration, ATTN: Lt. Col. Charles M. Barnes, USAF, DB-3, MSC, Houston, Texas 77058 (1)
- National Aeronautics and Space Administration, Manned Spacecraft Center, ATTN: Dr. B. D. Newsom, Mail Code DA, Houston, Texas 77058 (1)
- National Bureau of Standards, ATTN: Chief, Radiation Physics Division, Washington, D. C. 20234 (1)
- U. S. Public Health Service, Bureau of Radiological Health, Division of Biological Effects, 12720 Twinbrook Parkway, Rockville, Maryland 20852 (1)
- U. S. Public Health Service, Bureau of Radiological Health, Library, 12720 Twinbrook Parkway, Rockville, Maryland 20852 (1)
- U. S. Public Health Service, Northeastern Radiological Health Laboratory, 109 Holton Street, Winchester, Massachusetts 01890 (1)
- U. S. Public Health Service, Southeastern Radiological Health Laboratory, P. O. Box 61, Montgomery, Alabama 36101 (1)
- U. S. Public Health Service, Southwestern Radiological Health Laboratory, P. O. Box 15027, Las Vegas, Nevada 89114 (1)

## OTHER

- Argonne National Laboratory, Library Services Department, Report Section Bldg. 203, RM-CE-125, 9700 South Cass Avenue, Argonne, Illinois 60440 (1)
- Dr. Donald G. Baker, Radiobiology Department, Zellerbach Saroni Tumor Institute, 1600 Divisadero Street, San Francisco, California 94115 (1)
- Dr. J. T. Brennan, Radiology Department, University of Pennsylvania, 3400 Spruce Street, Philadelphia, Pennsylvania 19104 (1)
- Brookhaven National Laboratory, Information Division, ATTN: Research Library, Upton, Long Island, New York 11973 (2)
- Dr. J. S. Burkle, Director of Nuclear Medicine, York Hospital, York, Pennsylvania 17403 (1)
- Director, Radiobiology Laboratory, University of California, Davis, California 95616 (1)
- University of California, Lawrence Radiation Laboratory, Library, Bldg. 50, Room 134, Berkeley, California 94720 (1)
- University of California, Lawrence Radiation Laboratory, Technical Information Division Library L-3, P. O. Box 808, Livermore, California 94551 (2)
- University of California, Laboratory of Nuclear Medicine and Radiation Biology, Library, 900 Veteran Avenue, Los Angeles, California 90024 (1)
- Cdr. William H. Chapman, USN (Ret.), Bio-Medical Division L-523, Lawrence Radiation Laboratory, University of California, P. O. Box 808, Livermore, California 94551 (1)
- Director, Collaborative Radiological Health Laboratory, Colorado State University, Fort Collins, Colorado 80521 (1)
- Dr. L. W. Davis, Radiology Department, University of Pennsylvania, 3400 Spruce Street, Philadelphia, Pennsylvania 19104 (1)
- Professor Merrill Eisenbud, New York University, Tuxedo, New York 10987 (1)
- Dr. T. C. Evans, Radiation Research Laboratory, College of Medicine, University of Iowa, Iowa City, Iowa 52240 (1)
- Dr. Arnold Feldman, Institute of Radiology, School of Medicine, Washington University, 510 South Kingshighway, St. Louis, Missouri 63110 (1)
- Mr. Orin Gelderloos, Division of Literature, University of Michigan, Dearborn Campus, Dearborn, Michigan 48124 (1)
- General Dynamics/Fort Worth, ATTN: Librarian, P. O. Box 748, Fort Worth, Texas 76101 (1)
- Gulf General Atomic Incorporated, ATTN: Library, P. O. Box 608, San Diego, California 92112 (1)
- IIT Research Institute, ATTN: Document Library, 10 West 35th Street, Chicago, Illinois 60616 (1)
- Dr. R. F. Kallman, Department of Radiology, Stanford University, Palo Alto, California 94305 (1)
- Dr. L. S. Kelly, Donner Laboratory, University of California at Berkeley, Berkeley, California 94720 (1)
- Los Alamos Scientific Laboratory, ATTN: Report Librarian, P. O. Box 1663, Los Alamos, New Mexico 87544 (1)
- Director, Nuclear Science Center, Louisiana State University, Baton Rouge, Louisiana 70803 (2)
- Lovelace Foundation for Medical Education and Research, Document Library, 5200 Gibson Blvd., S. E., Albuquerque, New Mexico 87108 (1)
- Dr. Ross A. McFarland, Guggenheim Professor of Aerospace Health and Safety, Harvard School of Public Health, 665 Huntington Avenue, Boston, Massachusetts 02115 (1)
- Dr. J. I. Marcum, Rand Corporation, 1700 Main Street, Santa Monica, California 90401 (1)
- Massachusetts Institute of Technology, M. I. T. Libraries, Technical Reports, Room 14 E-210, Cambridge, Massachusetts 02139 (1)



## OTHER (continued)

- Dr. Charles W. Mays, Physics Group Leader, Radiobiology Division, University of Utah, Salt Lake City, Utah 84112 (1)
- Ohio State University, Nuclear Reactor Laboratory, 1298 Kinnear Road, Columbus, Ohio 43212 (1)
- Dr. Harvey M. Patt, Laboratory of Radiobiology, University of California, San Francisco Medical Center, San Francisco, California 94122 (1)
- Purdue University, Nuclear Engineering Library, Lafayette, Indiana 47907 (1)
- Dr. S. M. Reichard, Director, Division of Radiobiology, Medical College of Georgia, Augusta, Georgia 30902 (1)
- University of Rochester, Atomic Energy Project Library, P. O. Box 287, Station 3, Rochester, New York 14620 (1)
- Dr. H. H. Rossi, 630 West 168th Street, New York, N. Y. 10032 (1)
- Dr. Eugene L. Saenger, Director, Radioisotope Laboratory, Cincinnati General Hospital, Cincinnati, Ohio 45229 (1)
- Sandia Corporation Library, P. O. Box 5800, Albuquerque, New Mexico 87115 (1)
- Scientific Committee on the Effects of Atomic Radiation, ATTN: Library, United Nations Room 3267, United Nations Plaza, New York, N. Y. 10017 (1)
- Scope Publications, Franklin Station, P. O. Box 7407, Washington, D. C. 20004 (1)
- Dr. Arthur R. Tamplin, Biophysicist, Information Integration Group, University of California, Lawrence Radiation Laboratory, L-612, Livermore, California 94550 (1)
- Texas A. and M. University, Radiation Biology Laboratory, Texas Engineering Experiment Station, College Station, Texas 77840 (2)
- Texas Nuclear Corporation, ATTN: Director of Research, Box 9267 Allandale Station, Austin, Texas 78756 (1)
- Western Reserve University, Department of Radiology, Division of Radiation Biology, Cleveland, Ohio 44106 (1)
- Mr. Lionel Zamore, 601 Brightwater Court, Brooklyn, New York 11235 (1)

## FOREIGN

- International Atomic Energy Agency, Kärntnerring 11, Vienna I, 1010, Austria (1)
- European Atomic Energy Community, C. E. E. A., Library, 51 rue Belliard, Brussels 4, Belgium (1)
- Dr. L. G. Lajtha, Paterson Laboratories, Christie Hospital and Holt Radium Institute, Manchester, England (1)
- Dr. L. F. Lamerton, Biophysics Department, Institute of Cancer Research, Surrey Branch, Belmont, Sutton, Surrey, England (1)
- National Lending Library for Science and Technology, Boston Spa, Yorkshire, England (1)
- Directorate of Medical and Health Services, FAF (Federal Armed Forces), Bonn, Ermekeilstrasse 27, West Germany (1)
- Abteilung für Strahlenbiologie im Institut für Biophysik der Universität Bonn, 53 Bonn-Venusberg, Annaberger Weg 15, Federal Republic of Germany (2)
- Prof. Dr. H. Langendorff, Direktor des Radiologischen Instituts der Universität, 78 Freiburg im Breisgau, Albertstrasse 23, Germany (1)
- Priv.-Doz. Dr. O. Messerschmidt, Radiologisches Institut der Universität, 78 Freiburg im Breisgau, Albertstrasse 23, Germany (1)
- Dr. Helmut Mitschrich, Akademie des Sanitäts- und Gesundheitswesens der Bundeswehr, Spezialstab ATV, 8 München, Schwere Reiterstrasse 4, Germany (2)
- Prof. Dr. F. Wachsmann, Gesellschaft für Strahlenforschung m.b.H., 8042 Neuherberg bei München, Institut für Strahlenschutz, Ingolstadter Landstrasse 1, München, Germany (1)
- Col. Joachim Emde, Direktor, Spezialstab ATV, ABC- und Selbstschuttschule, 8972 Sonthofen 2/Allgäu, Berghoferstrasse 17, West Germany (1)
- Dr. M. Feldman, Section of Cell Biology, The Weizmann Institute of Science, Rehovoth, Israel (1)
- Dr. G. W. Barendsen, Radiobiological Institute TNO, Rijswijk, Netherlands (1)
- Puerto Rico Nuclear Center, ATTN: Reading Room, College Station, Mayaguez, Puerto Rico 00708 (2)
- Dr. H. Cottier, Pathological Institut der Universität, Bern, Switzerland (1)

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R & D

*Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified*

1. ORIGINATING ACTIVITY (Corporate author)

Armed Forces Radiobiology Research Institute  
Defense Atomic Support Agency  
Bethesda, Maryland 20014

2a. REPORT SECURITY CLASSIFICATION

UNCLASSIFIED

2b. GROUP

N/A

3. REPORT TITLE

ALTERATION IN GRAFT VERSUS HOST REACTIVITY AND IN HEMATOPOIETIC STEM CELLS OF SPLEEN CELL INOCULA FROM DONOR MICE PRETREATED WITH PERTUSSIS ANTIGEN

4. DESCRIPTIVE NOTES (Type of report and inclusive dates)

5. AUTHOR(S) (First name, middle initial, last name)

E. A. Elkman and R. T. Bowser

6. REPORT DATE

May 1971

7a. TOTAL NO. OF PAGES

30

7b. NO. OF REFS

52

8a. CONTRACT OR GRANT NO.

b. PROJECT NO. NWER XAXM

c. Task and Subtask C 903

d. Work Unit 01

9a. ORIGINATOR'S REPORT NUMBER(S)

AFRRI SR71-7

9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)

10. DISTRIBUTION STATEMENT

Approved for public release; distribution unlimited

11. SUPPLEMENTARY NOTES

12. SPONSORING MILITARY ACTIVITY

Director  
Defense Atomic Support Agency  
Washington, D. C. 20305

13. ABSTRACT

The graft versus host reactivity (GVHR) and hematopoietic stem cells of spleen cell inocula were assayed 3 days after pretreatment of donor mice with heated or unheated pertussis vaccine. Unheated vaccine reduced GVHR to 17 percent of control reactivity, measured by the Simonsen spleen weight assay, and increased 2.2-fold the proportion of hematopoietic stem cells, assayed as colony forming units (CFU). Heated vaccine, which did not produce the lymphocytosis induced in donors by unheated vaccine, reduced GVHR to 38 percent of control reactivity, and increased the proportion of CFU 3.8-fold. Alternative mechanisms for these effects are discussed.

DD FORM 1473 (PAGE 1)

1 NOV 65

S/N 0101-807-6801

UNCLASSIFIED

Security Classification